

FORM PTO-1599 (Modified) REV 11-98		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER <b>147-201P</b>	
<b>TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371</b>				U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR <b>09/555971</b>	
INTERNATIONAL APPLICATION NO. <b>PCT/EP98/07909</b>		INTERNATIONAL FILING DATE <b>4 December 1998</b>		PRIORITY DATE CLAIMED <b>5 December 1997 and 12 December 1997</b>	
TITLE OF INVENTION <b>METHOD FOR IDENTIFYING NUCLEIC ACIDS BY ELECTRO-SPRAY MASS SPECTROMETRY</b>					
APPLICANT(S) FOR DO/EO/US <b>GUT, Ivo Glynne; BERLIN, Kurt; and LEHRACH, Hana</b>					
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:					
<ol style="list-style-type: none"> <li>1. <input checked="" type="checkbox"/> This is a <b>FIRST</b> submission of items concerning a filing under 35 U.S.C. 371.</li> <li>2. <input type="checkbox"/> This is a <b>SECOND</b> or <b>SUBSEQUENT</b> submission of items concerning a filing under 35 U.S.C. 371.</li> <li>3. <input checked="" type="checkbox"/> This is an express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).</li> <li>4. <input checked="" type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.</li> <li>5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371 (c) (2))             <ol style="list-style-type: none"> <li>a. <input checked="" type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau).</li> <li>b. <input checked="" type="checkbox"/> has been transmitted by the International Bureau.</li> <li>c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US).</li> </ol> </li> <li>6. <input type="checkbox"/> A translation of the International Application into English (35 U.S.C. 371 (c)(2)).</li> <li>7. <input checked="" type="checkbox"/> A copy of the International Search Report (PCT/ISA/210).</li> <li>8. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))             <ol style="list-style-type: none"> <li>a. <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau).</li> <li>b. <input type="checkbox"/> have been transmitted by the International Bureau.</li> <li>c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired.</li> <li>d. <input checked="" type="checkbox"/> have not been made and will not be made.</li> </ol> </li> <li>9. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).</li> <li>10. <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).</li> <li>11. <input type="checkbox"/> A copy of the International Preliminary Examination Report (PCT/IPEA/409).</li> <li>12. <input type="checkbox"/> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).</li> </ol>					
<b>Items 13 to 20 below concern document(s) or information included:</b> <ol style="list-style-type: none"> <li>13. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98.</li> <li>14. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.</li> <li>15. <input type="checkbox"/> A <b>FIRST</b> preliminary amendment.</li> <li>16. <input type="checkbox"/> A <b>SECOND</b> or <b>SUBSEQUENT</b> preliminary amendment.</li> <li>17. <input type="checkbox"/> A substitute specification.</li> <li>18. <input type="checkbox"/> A change of power of attorney and/or address letter.</li> <li>19. <input checked="" type="checkbox"/> Certificate of Mailing by Express Mail</li> <li>20. <input checked="" type="checkbox"/> Other items or information:</li> </ol>					
<b>Six (6) sheets of drawings</b> <div style="border: 1px solid black; height: 100px; width: 100%;"></div>					

U.S. APPLICATION NO. (IF KNOWN) SEE OTHER

INTERNATIONAL APPLICATION NO.

ATTORNEY'S DOCKET NUMBER

097555971

PCT/EP98/07909

147-201P

21. The following fees are submitted:

**BASIC NATIONAL FEE ( 37 CFR 1.492 (a) (1) - (5) ) :**

- ☐ Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2) paid to USPTO and International Search Report not prepared by the EPO or JPO ..... \$970.00
- ☒ International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO ..... \$840.00
- ☐ International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO ..... \$690.00
- ☐ International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4) ..... \$670.00
- ☐ International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4) ..... \$96.00

**ENTER APPROPRIATE BASIC FEE AMOUNT =**

\$840.00

Surcharge of \$130.00 for furnishing the oath or declaration later than months from the earliest claimed priority date (37 CFR 1.492 (e)). ☐ 20 ☒ 30

\$130.00

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	21 - 20 =	1	x \$18.00
Independent claims	1 - 3 =	0	x \$78.00

\$18.00

\$0.00

Multiple Dependent Claims (check if applicable) ☒

\$260.00

**TOTAL OF ABOVE CALCULATIONS =**

\$1,248.00

Reduction of 1/2 for filing by small entity, if applicable. Verified Small Entity Statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28) (check if applicable). ☐

\$0.00

**SUBTOTAL =**

\$1,248.00

Processing fee of \$130.00 for furnishing the English translation later than months from the earliest claimed priority date (37 CFR 1.492 (f)). ☐ 20 ☒ 30

\$130.00

**TOTAL NATIONAL FEE =**

\$1,378.00

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable). ☐

\$0.00

**TOTAL FEES ENCLOSED =**

\$1,378.00

Amount to be refunded	\$
charged	\$

☒ A check in the amount of **\$1,378.00** to cover the above fees is enclosed.

☐ Please charge my Deposit Account No. \_\_\_\_\_ in the amount of \_\_\_\_\_ to cover the above fees.  
A duplicate copy of this sheet is enclosed.

☒ The Commissioner is hereby authorized to charge any fees which may be required, or credit any overpayment to Deposit Account No. **02-2448** A duplicate copy of this sheet is enclosed.

**NOTE:** Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive 37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

**BIRCH, STEWART, KOLASCH & BIRCH, LLP**  
P.O. Box 747  
Falls Church, VA 22040-0747

SIGNATURE

Leonard R. Svensson

NAME

30,330

REGISTRATION NUMBER

5 June 2000

DATE

**STATEMENT CLAIMING SMALL ENTITY STATUS**  
**(37 CFR 1.9(f) & 1.27(d))--NONPROFIT ORGANIZATION**

Docket No. \_\_\_\_\_

Applicant, Patentee, or Identifier: Ivo Glynne GUT et al.  
 Application No.: 09/555,971  
 Application Filed:  
 International Application No.: PCT/EP98/07909  
 International Filing Date: December 4, 1998  
 Title: Method for Identifying Nucleic Acids by Electro-Spray Mass Spectrometry

I hereby state that I am an official empowered to act on behalf of the nonprofit organization identified below:

NAME OF NONPROFIT ORGANIZATION: **Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V.**  
 ADDRESS OF NONPROFIT ORGANIZATION: **Berlin, Germany**

TYPE OF NONPROFIT ORGANIZATION:

- ☐ UNIVERSITY OR OTHER INSTITUTION OF HIGHER EDUCATION  
☐ TAX EXEMPT UNDER INTERNAL REVENUE SERVICE CODE (26 U.S.C. 501(a) and 501(c)(3))  
☐ NONPROFIT SCIENTIFIC OR EDUCATIONAL UNDER STATUTE OF STATE OF THE UNITED STATES OF AMERICA  
 (NAME OF STATE \_\_\_\_\_)  
 (CITATION OF STATUTE \_\_\_\_\_)  
☒ WOULD QUALIFY AS TAX EXEMPT UNDER INTERNAL REVENUE SERVICE CODE (26 U.S.C. 501(a) and 501(c)(3))  
 IF LOCATED IN THE UNITED STATES OF AMERICA  
☐ WOULD QUALIFY AS NONPROFIT SCIENTIFIC OR EDUCATIONAL UNDER STATUTE OF STATE OF THE UNITED  
 STATES OF AMERICA IF LOCATED IN THE UNITED STATES OF AMERICA  
 (NAME OF STATE \_\_\_\_\_)  
 (CITATION OF STATUTE \_\_\_\_\_)

I hereby state that the nonprofit organization identified above qualifies as a nonprofit organization as defined in 37 CFR 1.9(e) for purposes of paying reduced fees to the United States Patent and Trademark Office regarding the invention described in:

- ☐ the specification filed herewith with title as listed above.  
☒ the application identified above.  
☐ the patent identified above.

I hereby state that rights under contract or law have been conveyed to and remain with the nonprofit organization regarding the above identified invention. If the rights held by the nonprofit organization are not exclusive, each individual, concern, or organization having rights in the invention must file separate statements as to their status as small entities and that no rights to the invention are held by any person, other than the inventor, who would not qualify as an independent inventor under 37 CFR 1.9(c) if that person made the invention, or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or a nonprofit organization under 37 CFR 1.9(e).

Each person, concern, or organization having any rights in the invention is listed below:

- ☐ no such person, concern, or organization exists.  
☒ each such person, concern, or organization is listed below.

**Epigenomics GmbH**  
**Kastanienallee 24**  
**10435 Berlin, Germany**

(small business concern)

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b))

NAME OF PERSON SIGNING: Christa Herzog  
 TITLE IN ORGANIZATION OF PERSON SIGNING: Head of patent department  
 ADDRESS OF PERSON SIGNING: Hofgartenstr. 8, 80539 München

SIGNATURE: Christa Herzog DATE: 5.10.2000  
 Christa Herzog

09555971-111100

Max-Planck-Gesellschaft zur  
Förderung der Wissenschaften e.V.  
Our. ref.: B 3366 PCT

### Method for the identification of nucleic acid by means of electrospray mass spectrometry

The present invention relates to a method for detecting a nucleotide sequence in a nucleic acid molecule by means of predetermined probes with different masses each using electrospray mass spectrometry. Advantageously, the method of the invention allows at the same time for the characterisation of a variety of unknown nucleic acid molecules with a set of different probes. Moreover, this invention relates to a kit containing the probes and/or a probe support, optionally with nucleic acid molecules attached thereto.

The exact characterisation of nucleic acids is very time-consuming and expensive. It is possible to characterise an unknown DNA by sequencing it. This is the most precise way to analyse DNA. Sequencing of DNA, however, is very time-consuming and only necessary if the whole sequence is of interest. Only very short DNA segments (<1000 nucleobases) may be sequenced in one working process. If DNA fragments of more than 1000 nucleobases in length are to be sequenced on a large scale, it is necessary to subdivide the DNA, which makes the method more expensive.

Many statements, however, can also be made with a lower resolution. Yet, the methods that have been described up to now have the disadvantage that it might be necessary to use radioactivity and that only one probe can be used in one analysis. Such a method according to the state of the art, for instance, involves searching for partial information with an array of different target DNAs. An array of many thousands of target DNAs may be immobilized on a solid phase and subsequently, all target DNAs may be analysed together for the presence of a sequence by means of a probe (nucleic acid with a complementary sequence)<sup>1,2</sup>. A match of the target DNA with the probe can be proven by hybridizing the two nucleic acids. Probes may be any nucleic acid sequences of different lengths. There are different methods for selecting optimal libraries of probe sequences that overlap only to a very small extent<sup>3,4</sup>. It is also possible to compile the probe sequences purposefully in order to find certain target DNA sequences. Oligofingerprinting is an approach which uses this

technology. A library of target DNAs is scanned with short nucleic acid probes, with the probes mostly having a length of only 8-12 bases. One probe each is hybridized at once to a target DNA library that is immobilized on a nylon membrane. The probe is radioactively labelled and the hybridization is estimated according to the localisation of the radioactivity. For scanning an immobilized DNA array, fluorescently marked probes have also been used<sup>6</sup>. A similar method has been used for multiplexing the sequencing of DNA<sup>6,7</sup>. Use is made of various vector systems for cloning target sequences. One clone each is pooled by each cloning vector, the sequencing reaction is carried out and the fragments are separated on the gel and the gel is blotted onto a nylon membrane. The different sequences of the cloning system are subsequently hybridized to the immobilized DNA, so that the sequence belonging to the respective cloning system is obtained. Hereby, the scanning of the cloning system can also be carried out by means of a probe which is detectable by mass spectrometry<sup>8</sup>.

Any molecules capable of interacting sequence-specifically with a target DNA can be used as probes. Oligodeoxyribonucleotides are most commonly used. For this purpose, however, any modification of nucleic acids, e.g. peptide nucleic acids (PNA)<sup>9,10</sup>, phosphorothioate oligonucleotides or methyl phosphonate oligonucleotides, are suitable. The specificity of a probe is extremely important. Phosphothioate oligonucleotides are not particularly preferred as their structure is modified by the sulphur atoms, which has a negative influence on the nature of the hybridization. This can be due to the fact that phosphothioate oligonucleotides are usually not synthesized free of diastereomers. In the past, there has been a similar purity problem with methyl phosphonate oligonucleotides, but these oligonucleotides are increasingly synthesized free of diastereomers. An essential difference between methyl phosphonate oligonucleotides and phosphorothioate oligonucleotides is the uncharged backbone of the former, which leads to a reduced hybridization dependency on buffer salts and, all in all, to a higher affinity due to reduced rejection. Peptide nucleic acids also have an uncharged backbone which, at the same time, drastically deviates in its chemical properties from the common sugar phosphate structure of the backbone of nucleic acids. The backbone of a PNA has an amide sequence instead of a sugar phosphate backbone of normal DNA. PNA hybridizes very well to a sequence complementary to DNA. The melting temperature of a PNA/DNA hybrid is higher than the one of the corresponding DNA/DNA hybrid and the hybridization dependency on buffer salts is, again, relatively small. Electrospray mass spectrometry (ESI) is a new, very efficient method for analysing biomolecules (Meng, C.K., Mann, M. und Fenn, J.B., *Z. Phys. D.* 1988, 10, 361-368; Siuzdak, G. *Proc. Natl. Acad. Sci. USA*, 1994, 91, 11290-11297). The basic features of this

method are that a solution of a mixture of biomolecules is conducted through a metallic needle to which a high voltage is applied (typically several kilovolts), into the vacuum of a mass spectrometer. Due to the electric instability and the decreasing pressure, the solution disperses in small drips, each carrying electric charge on their surfaces. In several skimming steps, the drips are transferred to the high vacuum of the mass spectrometer. Thereby, the solvent of the drips is gradually lost and the naked, multiply charged biomolecules are located in an unfragmented form in the high vacuum of the mass spectrometer. For these methods, there are several possibilities as regards the step which comprises the mass spectrometrical analysis. One possibility relates to the separation of molecules with different mass and charge by means of a magnet (sector, quadrupole); another one relates to the analysis by means of measuring the time of flight. Hereby, the beam potential is pulsed vertically to the ion beam and the size of the particles is determined according to the time of flight. With these two embodiments it is possible to achieve an isotope resolution in the mass range of several thousand daltons. With new nanospray sources excellent spectrums can be recorded using only a few 100 fmol material.

Up to now, the ESI has essentially been successful for the analysis of peptides (Siuzdak, G., Proc. Natl. Acad. Sci. USA, 1994, 91, 11290-11297) and in particular for the analysis of peptide libraries produced combinatorially it has proven to be successful (Metzger, J.W., Kepter, C., Wiesmuller, K.H. and Jung, G., Anal. Biochem., 1994, 219, 261-277; Winger, B.E. and Campana, J.E., Rapid Commun. Mass Spectrom., 1996, 10, 1811-1813). The analysis of nucleic acids and modified variants has so far not been taken into consideration very often (Reddy, D.M., Rieger, R.A., Torres, M.C., and Iden, C.R., Anal. Biochem. 1994, 220, 200-207; Potier, N., Van Dorsselaer, A., Cordier, Y. Roch, O. and Bischoff, R., Nucleic Acids Res., 1994, 22, 3895-3903; Greig, M. and Griffey, R.H., Rapid Commun. Mass Spectrom., 1995, 9, 97-102). Thereby, problems arose as for example the fact that nucleic acid molecules are polyanions, which triggers additional problems in the ionisation.

Combinatorial syntheses<sup>13</sup>, i.e. the production of substance libraries starting from a mixture of preliminary stages are conducted in a liquid as well as in a solid phase. It is mainly the combinatorial solid phase synthesis that has established itself at a very early stage since the separation of by-products is particularly easy in this case. Only the target bonds attached to supports are retained in a washing step and at the end of the synthesis, they are isolated by specifically cleaving a linker. This technique allows in a very simple way for the simultaneous synthesis of a variety of different linkings to a solid phase and thus for obtaining chemically "pure" substance libraries<sup>14,15,16</sup>. Consequently, chemistry can easily access the classes of linkings which are also synthesised to a solid phase in combinatorial, conventional syntheses

and thus, these are widely used. This mainly applies to peptide, nucleic acid and PNA libraries.

The synthesis of peptides is carried out by attaching the first N-protected amino acid (e.g. Boc) to a support, by a subsequent removal of the protection and reaction of the second amino acid to the free  $\text{NH}_2$  group of the first one. In a further "capping" step, non-reacted amino functions are subjected to a further reaction in the next synthesis cycle. The protecting group at the amino function of the second amino acid is removed and the next building block may be coupled to it. For the synthesis of peptide libraries a mixture of amino acids is used in one or more steps. The synthesis of PNA and PNA libraries is carried out accordingly.

Nucleic acid libraries are mostly obtained by means of solid phase synthesis with mixtures of various phosphoramidite nucleosides. This may be conducted on commercially available DNA synthesizers without modifications in the synthesis protocols. Furthermore, various studies on the combinatorial synthesis of PNA libraries have been published<sup>17,18</sup>. These studies deal with the structure of combinatorial sequences, i.e. the synthesis of PNAs in which single, specific bases of a sequence are replaced by degenerated bases and thus, an incidental sequence variation is obtained. The use of mass spectrometrical methods for analysing combinatorial libraries has also been described several times<sup>19,20,21,22</sup>.

There are various methods for immobilizing DNA. The best-known technique is the solid linking of a DNA, which has been functionalized with biotin, to a streptavidin-coated surface<sup>23</sup>. The affinity of this system corresponds to a covalent chemical bond without being one. In order to be able to covalently bind a target DNA to a chemically prepared surface, a certain functionality of the target DNA is required. DNA itself does not have any functionalization that would be suitable for this purpose. There are several possibilities for introducing an appropriate functionalization into a target DNA: two functionalizations that are easy to work with are primary, aliphatic amines and thiols. Such amines quantitatively react with N-hydroxy succinimide esters. Under suitable conditions, thiols quantitatively react with alkyl iodides. One difficulty is the introduction of such a functionalization into a DNA. The simplest way is to introduce it by means of a primer of a PCR. In said methods a 5'-modified primer ( $\text{NH}_2$  and  $\text{SH}$ ) and a bifunctional linker are used<sup>24,25,26</sup>.

As regards the immobilization on a surface, it is mainly its nature that is of essential importance. Systems that have been described so far mainly consist of silicon or metal (magnetic beads). A further method for binding a target DNA is based on the

use of a short recognition sequence (e.g. 20 bases) in the target DNA for the hybridization to a surface-immobilized oligonucleotide<sup>27</sup>. Enzymatic techniques for introducing chemically active positions into target DNA have also been described<sup>28</sup>. In this case, a 5'-NH<sub>2</sub> functionalization is carried out enzymatically at a target DNA.

As has been described before, there is a number of methods known in the state of the art, which are all geared to an exact analysis of nucleic acids. Normally, these techniques are either very time-consuming and/or expensive.

Thus, the technical problem underlying the present invention was to provide a fast and cost-efficient method for the identification of target nucleic acids.

This technical problem has been solved by providing the embodiments characterised in the claims.

Therefore, the present invention relates to a method for detecting a nucleotide sequence in a nucleic acid molecule comprising the following steps:

- (a) hybridization of nucleic acid molecules to a set of probes of different nucleobase sequences, wherein each probe has a mass that differs from the one of all the other probes;
- (b) separation of the probes that were not hybridized;
- (c) detachment of a specifically hybridized probe in a solvent;
- (d) analysis of the hybridized probes in a solution by means of electrospray mass spectrometry; and
- (e) determination of the nucleic acid molecules by means of the probes hybridized to them.

Preferably, the nucleic acid molecules are fixed to a support. In this case, it is preferred that the positions of the probes on the probe support allow for an allocation to the nucleic acid molecule hybridizing therewith. The term "probe support", as used according to the invention, means any form of support material.

The method according to the invention (cf. Figure 1) advantageously combines methods for analysing target nucleic acids (oligofingerprinting), and the mass spectrometrical analysis of nucleic acids and modified nucleic acids. Thereby, a large number of different probes is used which allow for the detection of one or more nucleotide sequences in one nucleic acid molecule. So far, it has not been possible to combine these two methods, because the differences in the mass of the probes



0955974940

washing are carried out at a lower temperature, e.g. at 50°C or 55°C and/or the amount of SSC is raised, for example, to 1 x or 2 x SSC.

The method of the invention also allows for the detection of several different sequences in one target DNA, wherein the different sequences may be complementary to different probes. At the very best, e.g. when using probes with overlapping sequences, the whole nucleotide sequence of a target sequence may be detected or clarified.

With the method of the invention it may first be determined whether a target nucleic acid was fixed to an immobilization support, which exhibits a sequence that can be hybridized to a probe under the conditions selected. If this is the case, the nucleic acid of the sample may be further analysed and characterised. As in the method of the invention, usually only a very small percentage of a sample has to be used for the analysis according to the invention, the nucleic acid that is not used may be further analysed with standard methods, e.g. sequencing methods.

The removal of the specifically hybridized probes in a solvent is preferably conducted by means of thermal denaturation or alkaline denaturation. In particular, aqueous solutions or mixtures of water and organic solvents are suitable solvents, provided the organic solvent can be mixed with water (e.g. water and acetonitrile). Preferably, the mixtures are used at a ratio of 1:1. As the hybridized molecules are to be denaturated, the selection of the mixture, in addition, depends on the kind of molecules that are to be hybridized.

The method according to the invention may also be carried out several times either simultaneously or consecutively, with hybridization conditions being varied. In this way, it is possible, for example, to find out at first, whether a target DNA exhibits a specific degree of homology to other target DNAs, before searching for specific sequences.

The allocation of the probe hybridized at a specific position on the probe support to the immobilized sample is preferably conducted by means of a data processing system which allocates the respective spectrum recorded at a position of the probe support to a target DNA located exactly at the same position. Preferably, the target nucleic acids are located in a specific order on the surface or on the protein support.

In a preferred embodiment of the method according to the invention, the nucleic acid molecules are transferred to the surface of a support before or after step (a). For this

purpose, use can be made of a "magnetic bead", for example, which is functionalized in such a way that a target DNA functionalized correspondingly can be attached to it, or of a functionalization, for example with a microtitre plate in the different wells of which target DNAs may be immobilized. In this way, hybridization steps involving probe libraries and subsequent washing steps may be simplified. Biotin-streptavidin,  $\text{NH}_2$ -N-hydroxysuccinimidyl ester or SH-alkylhalide may be used.

In another preferred embodiment of the method of the invention, immobilization of the nucleic acid molecules is carried out at the surface of a probe support through a  $\text{NH}_2$ -, an epoxy- or a SH-function, through a protein-substrate, protein-protein or a protein-nucleic acid interaction by means of coating the surface of the probe support with a silicate or silane, or through an interaction of two hydrophobic building blocks.

If the probe support is coated with gold, coupling of the target DNA may take place by means of SH- or  $\text{NH}_2$ -functions introduced during the molecular biological preparation of the target DNA. There also is the reverse possibility of attaching a correspondingly modified DNA to functionalized gold particles. The company Nanoprobes Inc., Stony Brook, NY, for instance, sells gold-nanoparticles coupled with streptavidin or amino functions. By means of an amino functionalization, a subsequent coupling of the target DNA with a glass surface may be achieved. The target DNA is attached to a bifunctional linker (e.g. SIAB, Pierce Chemical, Rockford, IL, USA) through an SH-functionalization. Another possibility is to coat the surface directly with trimethoxy-3-amino propylsilane. Then, a target DNA may be coupled with the amino function through a bifunctional linker as described before.

In a particularly preferred embodiment, the protein-substrate interaction is a biotin-streptavidin or an antibody-antigen bond. Many probe supports usually consist of material to which neither proteins nor DNA molecules may be immobilized without further modification. One possibility of immobilizing is to coat the surface with gold since SH-functions, for example, may be coupled with it. For coupling, use can be made of bifunctional linkers having an SH-function and another function which is adapted to the functionalization of the target DNA. If the target DNA is, for example, biotin-functionalized, it should be possible to couple the linker with streptavidin. If the target DNA is  $\text{NH}_2$ -functionalized, the linker may have an N-hydroxysuccinimidyl ester-function.

In another particularly preferred embodiment, the protein-nucleic acid interaction is an attachment of the nucleic acid to Gene32, a protein which binds single-stranded DNAs in a way non-specific to sequences.

In another preferred embodiment of the method of the invention, the probes used are nucleic acids having a mass tag. According to this embodiment, the probes may also have several tags which may be located at different positions, e.g. at the 5' and the 3' end. The versatility and sensitivity of the method of the invention may be increased considerably by combining the amount and the localisation of mass tags, optionally in combination with charge tags.

In a particularly preferred embodiment the mass tags are at the same time charge tags, while in another particularly preferred embodiment, the nucleic acids additionally have a charge tag (cf. Figure 2).

Charge tagging may be conducted according to the method of Gut et al.<sup>11,12</sup>. An amino-functionalized substrate (1 mM) is put in trimethylamine/CO<sub>2</sub> buffer (pH = 8.5, 200 mM) on ice at 0°C with 1 % ω-trimethylammoniumhexanoic acid-N-hydroxy-succinimidylester (CT). After 30 minutes, the volatile buffer and the solvent in the vacuum are removed. The amino-functionalized substrate may, for example, be a combinatorially produced library of probes differing in their mass. By varying length and functionalization of the CT, the masses of the substrate library can be altered by a defined value (Fig. 2). As in the combinatorial synthesis, a probe library containing 64 probes differing in their mass in a range of 200 Da may be generated, the mass/charge tags increase the mass by units of 200 Da each, i.e. the first combinatorial synthesis preparation is generated with the smallest charge tag possible, the second with a mass/charge tag that has 200 Da more in weight, the third mass/charge tag that has another 200 Da more in weight, etc. In theory, this range can be extended at will as long as the mass spectrometer used is able to eliminate the difference between the two probes with neighbouring masses and as long as the synthesis seems to be feasible in practice. For probes with 10 nucleobases, a basic mass in the range of 2600 to 2800 Da is obtained. For mass spectrometers which can be bought at present, the range of mass that can be used and that has a sufficient accuracy as regards mass is below 4000 Da. Thus, seven ensembles of 64 probes can be used (all in all 448 probes).

The synthesis of peptides on an automated synthesizer takes place from the C-terminal end to the N-terminal end, the synthesis of nucleic acids from the 3' to the 5' end. There is the possibility of linking a primary amino function to one or to both ends in order to obtain mass shifting by means of one or two functionalizations. Alternatively, the masses of a library of combinatorially produced probes may also be altered by inserting some building blocks with a defined mass (e.g. amino acids in a combinatorial synthesis of PNAs) before inserting the combinatorial building blocks.

The first combinatorial synthesis starts directly at the support. For the second combinatorial synthesis, first of all two valines, for example, are coupled. Valine has a mass of 99 Da. With two valines the mass of the second combinatorial synthesis is changed by 198 Da compared to the first one. For the third combinatorial synthesis, first of all four valines are coupled, as a consequence, the mass of this ensemble is by 396 Da higher than the first one, etc. If a charge tag is necessary, it may be attached later on to the N-terminal end according to the above-mentioned method. It is also possible to couple the charge tags to the solid phase first and then to continue with the combinatorial synthesis (cf. for example Fig. 3).

Another possibility is to attach several fixed charges of the same polarity at the same time. Thus, the range of detection can be reduced. In a mass spectrometer, a molecule with two charges with the corresponding half of the mass is observed. This may be of advantage when the resolution of the mass spectrometer decreases drastically towards larger mass. It has been described before how to solidly link positive fixed charges to a probe. Negative fixed charges may have similar advantages. Positive and negative fixed charges can be generated with similar methods.

For the localisation, the amount and the combinatorics of the charge tags, the statements made in connection with the mass tags apply accordingly. Therefore, the sensitivity of an analysis may be improved by means of charge tagging at the 5' or 3' end of the library. This is particularly true if otherwise a neutral charge is generated by introducing alkylphosphonate groups into the non-randomized positions (cf. below). In addition, by means of mass tagging, which is inevitably related to charge tagging, the analysis of the fragments resulting from the cleavage at the phosphorothioate groups is facilitated. It is also possible to use phosphoroselenoates for achieving a preferred cleavage at certain positions.

In another preferred embodiment of the method of the invention, the probes are modified nucleic acid molecules.

In a particularly preferred embodiment, these modified nucleic acid molecules are PNAs, alkylated phosphorothioate nucleic acids or alkylphosphonate nucleic acids.

In another preferred embodiment, the probes of the method of the invention are generated by means of combinatorial solid phase synthesis. This, for example, can be carried out by means of the commonly used Boc solid phase synthesis which is also used commercially for the synthesis of PNA pure substances. In this case, all

four or more than one of the bases adenine, guanine, cytosine or thymine are incorporated next to each other in selected positions within a sequence of the probe DNA that is otherwise determined. This takes place by using not only one but several synthesis building blocks in one step of the solid phase synthesis. By varying several of these positions, a PNA library is created. After the PNA synthesis a charge tagging may be conducted at the free terminal amino function. This improves the dynamic range of the PNA analysis by means of ESI.

Therefore, in a preferred embodiment, the different building blocks for the solid phase synthesis are labelled in such a way that their mass or the mass of the probes synthesized from them can be differentiated in the mass spectrometer. This labelling is accomplished by introducing a different mass modification at the backbone in a way corresponding to the base. In this way, the synthesized probes obtain a mass that is specific to their sequences. If the probes now specifically bind to an immobilized target DNA, the information on the mass accessible in the mass spectrometrical experiment makes it possible to draw conclusions as to the sequences of the respective hybridized probe.

In another particularly preferred embodiment of the method of the invention, the building blocks are labelled with a methyl, ethyl, propyl, a branched or non-branched alkyl, a halogen-substituted branched or non-branched alkyl- or alkoxyalkyl, alkylaryl, arylalkyl, alkoxy or aryloxyalkyl group or with one of their deuterated or other isotopic variants. For, if building blocks are used the mass of which has not been modified, the molecular mass only makes it possible to draw conclusions as to the composition of the bases, but not as to the sequence, mass-labelled building blocks (i.e. PNA monomers substituted at the backbone) are used which are characteristic to any randomized position within the PNA library. As in this way the synthesis building block of a specific base has a different mass in position X than in position Y, it is possible to differentiate also different sequences with the same gross composition of the bases through the molecular mass. The respective substituents are selected by numeric values in such a way that a determined mass corresponds unequivocally to any possible sequence due to the described characteristics of the library synthesis.

The synthesis of nucleic acid probe libraries is conducted at the synthesizer by inserting mixtures of different nucleoside derivatives (usually phosphoramidites) at the positions to be randomized. The libraries resulting therefrom can also be used as probes in the above-described method. In order to differentiate the different sequences within the library, a specific bond breaking at the phosphodiester bonds located at a determined side of the randomized positions is to be carried out.

Therefore, in another preferred embodiment of the method of the invention, the probes have at least one modification in a defined position away from the randomized nucleotides which allow for the cleavage of the probe.

In a particularly preferred embodiment, this modification is the introduction of a phosphorothioate group and/or an RNA base and/or a phosphodiester bond.

If a probe contains three randomized positions, at least two such bond breakings are necessary (Fig. 6). As a result, three fragments are obtained which contain one randomized position each and which thus allow for a direct conclusion as to the variable base through its mass due to the composition otherwise known. The cleavage of the described bonds may also be conducted incompletely. This makes it possible to include larger fragments to secure the sequence information or to concretize it in the case of ambiguities. The specific bond breaking at the randomized positions is carried out by introducing phosphorothioate groups already during the synthesis of the library. These may first be hydroxyalkylated with hydroxyalkylhalides and then cleaved selectively under alkaline conditions. Alternatively, the samples may be set up in such a way that uracil is inserted next to a randomized position. The backbone can then be broken at the same position by means of uracil DNA glycosylase and subsequent alkaline treatment.

The above-described principle of mass tagging for certain components in determined positions may be used for various partial libraries which then may be united to a larger library.

Therefore, in another preferred embodiment, the probes are generated as partial libraries having different masses and/or charge tags. It is necessary to mass-label also the partial libraries in their syntheses so that specific analysed masses allow for conclusions as to that specific partial library. Usefully, this is carried out through natural amino acids which are easy to attach to a PNA library in a solid phase synthesis. By means of mass-labelling different partial libraries, it is possible to considerably extend the range of mass which is disposable for the analysis of the total library.

Moreover, the present invention relates to a kit containing the probes and/or parts of an immobilization system (probe supports), optionally with the nucleic acid molecules bound. (Coated) "magnetic beads" or a (coated) microtitre plate are suitable immobilization systems. As has been described before, the surface of the

immobilization system is pretreated and thus allows for the binding of the nucleic acids. Preferably, this pretreatment is carried out in a chemical way.

In this application, a number of publications are cited which are herewith incorporated into the specification by reference.

The Figures show:

#### Figure 1

Scheme of fingerprinting with mass spectrometrical scanning.

1. Combinatorially generated library of probes that can be differentiated by their mass.
2. The library is hybridized to an immobilized target DNA. After washing it intensively (to remove non-specifically bound probes), the correctly hybridized probes are separated from the target DNA.
3. This solution is analysed in an electrospray mass spectrometer. The partial sequences of the target DNA are determined by clearly allocating the mass to the sequences.

#### Figure 2

N-terminal mass/charge tagging.

A charge tag may be introduced at the 5' of a nucleic acid or a modified version of a nucleic acid, or at the N-terminal end of a PNA. The charge tag has an N-hydroxy-succinimidester function and a quaternary ammonium group. These two functionalizations are separated by one group  $R_1$ .  $R_1$  is used to vary the mass of the charge tags (coupling of the charge tags is carried out at a slightly alkaline pH (8.5) in an aqueous solution on ice. The reaction is completed within 30 minutes).

#### Figure 3

Principle of synthesis of PNA libraries using syntheses of mass-labelled syntheses that are carried out in parallel as well as mass-labelled building blocks L (randomized position). Various bases with different substituents each are used (as mass tags) in



the randomized positions at the backbone. The mass of the respective PNA molecule is allocated unequivocally to its sequence.

#### Figure 4

Calculated mass spectrum of a PNA library.

Two different solid phase syntheses (one of which is mass-labelled) with 32 different sequences each make up 64 different mass peaks, each of them being allocated to a specific sequence from a PNA library with 3 variable positions with four bases (A, C, G and T being inserted). The calculation is based on the substituents listed in Table 1. For the calculations the computer programme MASP (© Dr. Christoph Steinbeck) was used.

#### Figure 5

Possible synthesis components for the combinatorial Boc solid phase synthesis as are also used for the synthesis of PNA libraries that can be differentiated due to their mass. The components listed in Table 1 are shown.

#### Figure 6

Sequence analysis of probes by means of specific bond breakings.

The different sequences of a library of DNA probes can also be identified through mass spectrometry by means of specific bond breakings at the randomized positions ( $N = A, G, C$  or  $T$ ). A phosphorothioate function at which the DNA may be cleaved specifically (e.g. with idoethanol) is inserted during the solid phase synthesis. This is shown schematically without taking the exact chemical nature of the fragments into account. Since the sequence is known to a great extent, the mass of the fragments makes it possible to determine the total sequence without having to carry out a complete sequencing. The substituent  $R_1$  stands for any possible further DNA sequence,  $R_2$  for another possible sequence or  $-H$ .  $R_3$  stands for  $-OH$  (phosphodiester) or alkyl- (alkylphosphonate).

The examples explain the invention.

Example 1: Explanation of an embodiment of the whole method of the invention

A target DNA is generated with a function allowing for its immobilization at a surface. The target DNA is produced by means of PCR and in said PCR, a primer functionalized with biotin is used. For the immobilization, the surface is coated with streptavidin. A library of probes that can be differentiated due to their mass is hybridized to a target DNA in the above-described way. Complementary building blocks of the probe library bind to the respective target DNAs. Then, they are washed thoroughly to remove non-specifically bound probes. The specific probes are separated from the target DNA by means of heat denaturation. The solution is analysed in an electrospray mass spectrometer. Due to the mass found and the fact that the mass of the probes is distinguishable conclusions as to partial sequences in the target DNA can be drawn.

Example 2: Analysis of target DNA immobilized via a biotin-streptavidin bridge

A target DNA is attached to a streptavidin-coated magnetic bead by means of a biotin function. A library of probes distinguishable by mass is hybridized to target DNA. Non-specific hybridizations are washed off. By heating in an aqueous solution, the specific hybridized probes are detached. The aqueous solution is analysed in an electrospray mass spectrometer. Due to the mass found and the fact that the mass of the probes is distinguishable conclusions as to partial sequences in the target DNA can be drawn.

Example 3: Immobilization of target DNA

In a preferred variant of the method, the target DNA is attached to a bead consisting of an epoxy-functionalized acrylpolymer. Subsequently, the epoxy functions that have not reacted are deactivated with a surplus of an amine. The example is continued as described in Example 2.

Example 1: Explanation of an embodiment of the whole method of the invention

A target DNA is generated with a function allowing for its immobilization at a surface. The target DNA is produced by means of PCR and in said PCR, a primer functionalized with biotin is used. For the immobilization, the surface is coated with streptavidin. A library of probes that can be differentiated due to their mass is hybridized to a target DNA in the above-described way. Complementary building blocks of the probe library bind to the respective target DNAs. Then, they are washed thoroughly to remove non-specifically bound probes. The specific probes are separated from the target DNA by means of heat denaturation. The solution is analysed in an electrospray mass spectrometer. Due to the mass found and the fact that the mass of the probes is distinguishable conclusions as to partial sequences in the target DNA can be drawn.

Example 2: Analysis of target DNA immobilized via a biotin-streptavidin bridge

A target DNA is attached to a streptavidin-coated magnetic bead by means of a biotin function. A library of probes distinguishable by mass is hybridized to target DNA. Non-specific hybridizations are washed off. By heating in an aqueous solution, the specific hybridized probes are detached. The aqueous solution is analysed in an electrospray mass spectrometer. Due to the mass found and the fact that the mass of the probes is distinguishable conclusions as to partial sequences in the target DNA can be drawn.

Example 3: Immobilization of target DNA

In a preferred variant of the method, the target DNA is attached to a bead consisting of an epoxy-functionalized acrylpolymer. Subsequently, the epoxy functions that have not reacted are deactivated with a surplus of an amine. The example is continued as described in Example 2.

Example 4: Analysis of various target DNAs in a microtitre plate

In a microtitre plate 96 or 384 different biotin-functionalized target DNAs are immobilized in one well each. In each well, hybridization to a library of probes distinguishable by mass is carried out. The plate is washed in order to remove non-specifically bound hybridizations. The specific hybridizations are solved in the respective well by heating in an aqueous solution. The aqueous solution in each well is analysed in an electrospray mass spectrometer. Due to the mass found and the fact that the mass of the probes is distinguishable allow for conclusions as to partial sequences of the target DNA in the respective well.

Example 5: Synthesis of a PNA library with 64 different sequences

Figure 4 and Table 1 show an example of a PNA library with a clear relation between mass and sequence. In an otherwise random PNA, three positions are to be varied in the combinatorial solid phase synthesis. With 4 different bases this corresponds to 64 possible linkings. In the present example, two separate syntheses are conducted with the building blocks listed in Table 1. The second synthesis is additionally mass-labelled by adding two valine units. By each synthesis 32 compounds are obtained which can all be differentiated by their mass. Both partial libraries can be united to one library containing 64 different sequences, each one with a specific mass. The 64 peaks do not overlap, each peak corresponds to a specific sequence from a PNA library with three randomized base positions (4 bases each).

Example 6: Creation of a probe library with 448 probes

7 syntheses are carried out according to the method described in the latter example and subsequently, the mass/charge tags are coupled.

	mass range
1. 64 probes: charge tag- TCP <sub>1</sub> GAP <sub>2</sub> GAP <sub>3</sub> G	2600-2800 Da
2. 64 probes: charge tag+200 Da mass tag- TCP <sub>1</sub> AGP <sub>2</sub> GAP <sub>3</sub> G	2800-3000 Da
3. 64 probes: charge tag+400 Da mass tag- TCP <sub>1</sub> AGP <sub>2</sub> AGP <sub>3</sub> G	3000-3200 Da
4. 64 probes: charge tag+600 Da mass tag- TCP <sub>1</sub> AAP <sub>2</sub> AGP <sub>3</sub> G	3200-3400 Da
5. 64 probes: charge tag+800 Da mass tag- TCP <sub>1</sub> AAP <sub>2</sub> GAP <sub>3</sub> G	3400-3600 Da
6. 64 probes: charge tag+1000 Da mass tag- TCP <sub>1</sub> GAP <sub>2</sub> GAP <sub>3</sub> G	3600-3800 Da

7. 64 probes: charge tag+1200 Da mass tag- TCP<sub>1</sub>GAP<sub>2</sub>AGP<sub>3</sub>G 3800-4000 Da

In the above-mentioned synthesis series, the sixth synthesis serves as an internal control. Alternatively, the following synthesis may also be conducted:

6. 64 probes: charge tag+1000 Da mass tag- TCP<sub>1</sub>GGP<sub>2</sub>GAP<sub>3</sub>G 3600-3800 Da.

Example 7: Coating of a microtitre plate with protein

The surface of a microtitre plate is coated with Gene32, a protein that non-specifically binds single-stranded DNA. After coating the target with this protein, an array of target DNAs can be put onto it. If the array of the target DNAs consists of cDNA, these were primed with oligo-dT (e.g. dTTTTTTTTTTTTT) in this PCR. Oligo-dT interacts heavily with Gene32. The covalent coupling of the oligo-dT to Gene32 may be achieved by means of photocrosslinking with short UV light<sup>29,30</sup>. After this immobilization, a library of probes may be used for analysing the target DNA in the above-described way. Use may also be made of sequence-specific protein/DNA interaction (e.g. GCN4/AP1).

Example 8: Immobilization of target DNA at an epoxy-functionalized solid phase.

10 mg support (Eupergit C250 L, Röhm Pharma Polymere) are suspended in 1 ml immobilization buffer (1 M K<sub>2</sub>HPO<sub>4</sub>/KH<sub>2</sub>PO<sub>4</sub>, pH 7.5). 1.5 nmol of the target DNA that is to be immobilized is added and incubated for 24 hours at room temperature while shaking gently. The supernatant is removed and the epoxy functions which have not reacted are deactivated through treatment with 1 M glycine solution for 24 hours at room temperature. The supernatant is removed and the support is washed several times. In this way, the immobilized DNA can be stored for a longer time at -20°C.

Example 9: Selective hybridization with "charge tagged" PNAs

The immobilized target DNA is incubated for 15 min at 65°C with the library of "charge tagged" PNA probes (80 pmol in 20 µl/8 pmol immobilized DNA) in hybridization buffer (10 mM Tris/HCl), 5 mM NH<sub>4</sub>Cl, 15 % formamide) and subsequently, the probes that have not been hybridized are removed by washing

three times with hybridization buffer. Then, the dehybridization is carried out in 40% acetonitril at 80°C. An aliquot of this solution is lyophilized, put into water and analysed in a ESI mass spectrometer.

base	position 1	position 2	position 3
A	H	<i>i</i> Pr	H
T	H	Me	<i>i</i> PrOCH <sub>2</sub>
C	H	H	<i>i</i> PrOCH <sub>2</sub> *
G	H	<i>i</i> Bu	H*

Table 1: Substituents fixed at the PNA subunits for producing a mass-labelled library with an unequivocal mass/sequence.

Relation. \*: second synthesis, mass-labelled with 2 valine units. Figure 5 shows the corresponding synthesis building blocks.

## References

- <sup>1</sup> Hoheisel, J.D. and Lehrach, H. 1993. Use of reference libraries and hybridisation/fingerprint for relational genome analysis. *FEBS*. 325: 118-122.
- <sup>2</sup> Scholler, P., Karger, A.E., Meier-Ewert, S., Lehrach, H., Delius, H. and Hoheisel, J.D. 1995. Fine-mapping of shotgun template-libraries; an efficient strategy for the systematic sequencing of genomic DNA. *Nucleic Acids Res.* 23: 3842-3849.
- <sup>3</sup> Brenner, S. 1994. Methods for sorting polynucleotides using oligonucleotide tags. US Patent. 5,604,097.
- <sup>4</sup> Brenner, S. 1995. Minimally cross-hybridizing sets of oligonucleotide tags. US Patent. 5,635,400.
- <sup>5</sup> Guo, Z., Guilfoyle, R.A., Thiel, A.J., Wang, R. and Smith, L.M. 1994. Direct fluorescence analysis of genetic polymorphisms by hybridization with oligonucleotides arrays of glass supports. *Nucleic Acids Res.* 22: 5456-5465.
- <sup>6</sup> Church, G.M. and Kieffer-Higgins, S. 1988. Multiplex DNA sequencing. *Science*. 240: 185-188.
- <sup>7</sup> Church, G.M. and Kieffer-Higgins, S. 1990. Multiplex analysis of DNA. US Patent. 5,149,625.
- <sup>8</sup> Arlinghaus, H.F., Kwoka, M.N., Guo, X.-Q. and Jacobson, K.B. 1997. Multiplex DNA sequencing and diagnostics by hybridization with enriched stable isotope labels. *Anal. Chem.* 69: 1510-1517.
- <sup>9</sup> Nielsen, P.E., Buchardt, O., Egholm, M. and Berg, R.H. 1993. Peptide nucleic acids. US Patent. 5,539,082.

- 10 Buchardt, O., Egholm, M., Berg, R.H. and Nielsen, P.E. 1993. Peptide nucleic acids and their potential applications in biotechnology. *Trends in Biotechnology*, 11: 384-386.
- 11 Gut, I.G. and Beck, S. 1995. Method of nucleic acid analysis. Patent WO96/27681.
- 12 Gut, I.G., Jeffery, W.A., Pappin, D.J.C. and Beck, S. 1997. Analysis of DNA by 'charge tagging' and matrix-assisted laser desorption/ionization mass spectrometry. *Rapid Commun. Mass Spectrom.* 11: 43-50.
- 13 Lowe, G. 1995. Combinatorial Chemistry. *Chem. Soc. Rev.* 24: 309.
- 14 Jung, G., Früchtel, J.S. 1996. Organische Chemie an fester Phase. *Angew. Chem.* 108: 19-46.
- 15 Choong, I.C., Ellman, J.A. 1996. Solid-Phase Synthesis: Applications to Combinatorial Libraries. *Ann. Reports in Med. Chem.* 31: 309.
- 16 Pirrung, M.C. 1997. Spatially Addressable Combinatorial Libraries. *Chem. Rev.* 97: 473.
- 17 Cook, P.D., Kiely, J. and Sprankle, K. 1994. Peptide nucleic acid combinatorial libraries and improved methods of synthesis. US Patent. 5,539,083.
- 18 Nielsen, P.E. Peptide nucleic acids: a new dimension to peptide libraries and aptamers. *Methods in Enzymology.* 426-433.
- 19 Metzger, J.W., Stevanovic, S., Brünjes, J., Wiesmüller, K.-H., Jung, G. 1994. Electrospray Mass Spectrometry and Multiple Sequence Analysis of Synthetic Peptide Libraries. *Methods: A Companion to Methods in Enzymology* 6: 425-431.
- 20 Loo, J.A., DeJohn, D.E., Andrews, P.C. 1996. Application of Mass Spectrometry for Characterizing and Identifying Ligands from Combinatorial Libraries. *Ann. Reports in Med. Chem.* 31:319.



- 21 Pomerantz, S.C., McCloskey, J.A., Eaton, B.C., 1997. Deconvolution of Combinatorial Oligonucleotide Libraries by Electrospray Ionization Tandem Mass Spectrometry. *J. Am. Chem. Soc.* 119: 3861.
- 22 Carr, S.A., Benkovic, S.J., Winograd, N. 1996. Evaluation of Mass Spectrometric Methods Applicable to the Direct Analysis of Non-Peptide Bead-Bound Combinatorial Libraries. *Anal. Chem.* 68: 237.
- 23 Uhlen, M. et al. 1988, *Nucleic Acids Res.* 16, 3025-3038.
- 24 Chrisey, L.A., Lee, G.U. and O'Ferrall, C.E. 1996. Covalent attachment of synthetic DNA to self-assembled monolayer films. *Nucleic Acids Res.* 24: 3031-3039.
- 25 Timofeev, E.N., Kochetkova, S.V., Mirzabekov, A.D. and Florentiev, V.L. 1996. Regioselective immobilization of short oligonucleotides to acrylic copolymer gels. *Nucleic Acids Res.* 24: 3142-3148.
- 26 O'Donnell, M.J., Tang, K., Köster, H., Smith, C.L. and Cantor, C.R. 1997. High-density, covalent attachment of DNA to silicone wafers for analysis by MALDI-TOF mass spectrometry. *Anal. Chem.* 69: 2438-2443.
- 27 Cantor, C.R. 1995. Methods of preparing probe array by hybridization. US Patent. 5,631,134.
- 28 Bruick, R.K., Koppitz, M., Joyce, G.F. and Orgel, L.E. 1997. A simple procedure for constructing 5'-amino-terminated oligodeoxynucleotides in aqueous solution. *Nucleic Acids Res.* 25: 1309-1310.
- 29 Hockensmith, J.W., Kubasek, W.L., Vorachek, W.R. and von Hippel, P.H. 1993. Laser cross-linking of proteins to nucleic acids. *J. Bio. Chem.* 268: 15712-15720.
- 30 von Hippel, P.H. 1994. Protein-DNA recognition: New perspectives and underlying themes. *Science.* 263: 769-770.

# VOSSIUS & PARTNER

Patentanwälte

SIEBERTSTRASSE 4 · 81675 MÜNCHEN

TEL.: +49-89-4130 40 · FAX: +49-89-4130 41 11 · FAX (Marken-Trademarks): +49-89-41 30 44 00

## Translation of the amended claims

1. Method for detecting a nucleotide sequence in a nucleic acid molecule comprising the following steps:
  - (a) hybridization of nucleic acid molecules to a set of probes of different nucleobase sequences, wherein each probe has a mass that differs from the one of all the other probes;
  - (b) separation of the probes that were not hybridized;
  - (c) detachment of a specifically hybridized probe in a solvent;
  - (d) analysis of the hybridized probes in a solution by means of electrospray mass spectrometry; and
  - (e) determination of the nucleic acid molecules by means of the probes hybridized to them.
2. Method according to claim 1, wherein the nucleic acid molecules are immobilized at the surface of a support before or after step (a).
3. Method according to claim 2, wherein the immobilization of the nucleic acid molecules at the surface is carried out via an  $\text{NH}_2$ , epoxy or SH function by means of coating the surface of the probe supports with a silicate or silane, via a protein-substrate, protein-protein or a protein-nucleic acid interaction or via an interaction of two hydrophobic building blocks.
4. Method according to claim 3, wherein the protein-substrate interaction is a biotin-streptavidin bond or an antibody-antigen bond.
5. Method according to claim 3, wherein the protein-nucleic acid interaction is a Gene32-nucleic acid bond.
6. Method according to any one of claims 1 to 5, wherein the probes are nucleic acids having a mass tag.

7. Method according to claim 6, wherein the mass tag is at the same time a charge tag.
8. Method according to claim 6, wherein the nucleic acids moreover have a charge tag.
9. Method according to any one of claims 1 to 8, wherein the probes are modified nucleic acid molecules.
10. Method according to claim 9, wherein the modified nucleic acid molecules are PNAs, alkylated phosphorothioate nucleic acids or alkylphosphonate nucleic acids.
11. Method according to any one of claims 1 to 10, wherein the probes are generated by means of combinatorial solid phase synthesis.
12. Method according to claim 11, wherein different base building blocks are labelled in such a way that the probes synthesized therefrom can be differentiated in the mass spectrometer due to their mass.
13. Method according to claim 12, wherein the labelling is a methyl, ethyl, propyl, a branched or non-branched alkyl, a halogen substituted branched or non-branched alkyl, alkoxyalkyl, alkylaryl, arylalkyl, alkoxyaryl or aryloxyalkyl group or one of their deuterated or other isotopic variants.
14. Method according to any one of claims 10 to 13, wherein the probes have at least one modification in a defined position away from randomized nucleotides allowing for the cleavage of the probe.
15. Method according to claim 14, wherein modification means the introduction of a phosphorothioate group and/or an RNA base and/or a phosphotriester bond into the probe.


16. Method according to any one of claims 1 to 15, wherein the probes are generated as partial libraries having different mass and/or charge tags.
17. Method according to any one of claims 1 to 16, wherein the positions of the probes on the probe support allow for an allocation to the nucleic acid molecules hybridizing thereto.
18. Kit comprising
- (a) a set of probes as defined in any one of claims 6 to 16 and/or
  - (b) a probe support which has been pretreated and thus allows for the attachment of target DNAs and/or target DNAs that have already been attached.

## Abstract


The present invention relates to a method for detecting a nucleotide sequence in a nucleic acid molecule by means of predetermined probes with different mass using electrospray mass spectrometry. Advantageously, the method of the invention allows at the same time for the characterisation of a variety of unknown nucleic acid molecules by means of different probes. Moreover, the invention relates to a kit comprising the probes and/or a probe support, optionally with nucleic acid molecules attached thereto.

0555971-14400

## 1. Mass distribution of the probes



## 2. Hybridization



## 3. Mass distribution of the hybridized probes






Fig. 1

# N-terminal mass/charge tagging



R = e.g. alkyl, -CH<sub>3</sub>, -C<sub>2</sub>H<sub>5</sub>, -C<sub>3</sub>H<sub>7</sub>, -C<sub>4</sub>H<sub>9</sub> etc.



R<sub>2,4</sub> = e.g. alkyl, substituted alkyl

R<sub>5</sub> = e.g. nucleic acid, PNA, methyl phosphonate nucleic acid, phosphorothioate nucleic acid

Fig. 2





Fig. 3

B = adenine, cytosine, guanine, thymine or purine or pyrimidine derivatives or their deaza analogues

L(n) are various sets of substituents, selected specifically for each base, which are inserted in each synthesis step in order to obtain minimized peak overlaps in the MALDI-MS.





64 mass peaks corresponding to a specific PNA sequence;  
mass tagging

Fig. 4




Fig. 5




Fig. 6

# BIRCH, STEWART, KOLASCH & BIRCH, LLP

P.O. Box 747 • Falls Church, Virginia 22040-0747  
Telephone: (703) 205-8000 • Facsimile: (703) 205-8050

PLEASE NOTE:  
YOU MUST  
COMPLETE THE  
FOLLOWING

## COMBINED DECLARATION AND POWER OF ATTORNEY FOR PATENT AND DESIGN APPLICATIONS

As a below named inventor, I hereby declare that: my residence, post office address and citizenship are as stated next to my name; that I verily believe that I am the original, first and sole inventor (if only one inventor is named below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

Insert Title:

METHOD FOR IDENTIFYING NUCLEIC ACIDS BY ELECTRO-SPRAY MASS SPECTROMETRY

Fill in Appropriate  
Information -  
For Use Without  
Specification  
Attached:

the specification of which is attached hereto. If not attached hereto,

the specification was filed on June 5, 2000 \_\_\_\_\_ as

United States Application Number 09/555,971 \_\_\_\_\_;

\_\_\_\_\_ (if applicable) and/or

\_\_\_\_\_ as PCT

International Application Number PCT/EP98/07909 \_\_\_\_\_; and was

amended under PCT Article 19 on \_\_\_\_\_ (if applicable)

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I do not know and do not believe the same was ever known or used in the United States of America before my or our invention thereof, or patented or described in any printed publication in any country before my or our invention thereof or more than one year prior to this application, that the same was not in public use or on sale in the United States of America more than one year prior to this application, that the invention has not been patented or made the subject of an inventor's certificate issued before the date of this application in any country foreign to the United States of America on an application filed by me or my legal representative or assigns more than twelve months (six months for designs) prior to this application, and that no application for patent or inventor's certificate on this invention has been filed in any country foreign to the United States of America prior to this application by me or my legal representative or assigns, except as follows.

I hereby claim foreign priority benefits under Title 35, United States Code, §119(a)-(d) of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

### Prior Foreign Application(s)

### Priority Claimed

Insert Priority  
Information:  
(if appropriate)

97121470.5

European Patent

December 5, 1997

☒ ☐

(Number)

(Country)

(Month/Day/Year Filed)

Yes No

97121983.7

European Patent

December 12, 1997

☒ ☐

(Number)

(Country)

(Month/Day/Year Filed)

Yes No

\_\_\_\_\_  
(Number)

\_\_\_\_\_  
(Country)

\_\_\_\_\_  
(Month/Day/Year Filed)

☐ ☐

\_\_\_\_\_  
(Number)

\_\_\_\_\_  
(Country)

\_\_\_\_\_  
(Month/Day/Year Filed)

☐ ☐

\_\_\_\_\_  
(Number)

\_\_\_\_\_  
(Country)

\_\_\_\_\_  
(Month/Day/Year Filed)

☐ ☐

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional applications(s) listed below.

Insert Provisional  
Application(s):  
(if any)

\_\_\_\_\_  
(Application Number)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Application Number)

\_\_\_\_\_  
(Filing Date)

All Foreign Applications, if any, for any Patent or Inventor's Certificate Filed More than 12 Months (6 Months for Designs) Prior to the Filing Date of This Application:

Country

Application Number

Date of Filing (Month/Day/Year)

Insert Requested  
Information:  
(if appropriate)

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

I hereby claim the benefit under Title 35, United States Code, §120 of any United States and/or PCT application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States and/or PCT application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose information which is material to the patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

Insert Prior U.S.  
Application(s):  
(if any)

\_\_\_\_\_  
(Application Number)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Status - patented, pending, abandoned)

\_\_\_\_\_  
(Application Number)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Status - patented, pending, abandoned)

I hereby appoint the following attorneys to prosecute this application and/or an international application based on this application and to transact all business in the Patent and Trademark Office connected therewith and in connection with the resulting patent based on instructions received from the entity who first sent the application papers to the attorneys identified below, unless the inventor(s) or assignee provides said attorneys with a written notice to the contrary:

Raymond C. Stewart  
Joseph A. Kolasch  
Bernard L. Sweeney  
Charles Gorenstein  
Leonard R. Svensson  
Andrew D. Meikle  
Joe McKinney Muncy  
John W. Bailey  
Gary D. Yacura

(Reg. No. 21,066)  
(Reg. No. 22,463)  
(Reg. No. 24,448)  
(Reg. No. 29,271)  
(Reg. No. 30,330)  
(Reg. No. 32,868)  
(Reg. No. 32,334)  
(Reg. No. 32,881)  
(Reg. No. 35,416)

Terrell C. Birch  
James M. Slattery  
Michael K. Mutter  
Gerald M. Murphy, Jr.  
Terry L. Clark  
Marc S. Weiner  
Donald J. Daley  
John A. Castellano

(Reg. No. 19,382)  
(Reg. No. 28,380)  
(Reg. No. 29,680)  
(Reg. No. 29,977)  
(Reg. No. 32,644)  
(Reg. No. 32,811)  
(Reg. No. 34,313)  
(Reg. No. 35,094)

Send Correspondence to:

**BIRCH, STEWART, KOLASCH & BIRCH, LLP**  
P.O. Box 747 • Falls Church, Virginia 22040-0747  
Telephone: (703) 205-8000 • Facsimile: (703) 205-8050

or

Customer No. 2292

PLEASE NOTE

YOU MUST  
COMPLETE  
THE  
FOLLOWING:

Full Name of First  
Inventor  
Insert Date That  
Document is Signed

Insert Residence  
Insert Citizenship

Insert Post Office  
Address

Full Name of Second  
Inventor, if any  
see above

Full Name of Third  
Inventor, if any  
see above

Full Name of Fourth  
Inventor, if any  
see above

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

GIVEN NAME/FAMILY NAME Ivo Glynne GUT		INVENTOR'S SIGNATURE <i>Ivo Glynne Gut</i>	DATE* 3. October 2020
Residence (City, State & Country) Paris, France		CITIZENSHIP Swiss / British	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country) 18 rue du Moulin Vert, 75014 Paris, France			
GIVEN NAME/FAMILY NAME Kurt BERLIN		INVENTOR'S SIGNATURE <i>Kurt Berlin</i>	DATE*
Residence (City, State & Country) Stahnsdorf, Germany		CITIZENSHIP German	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country) Marienkaferweg 4, 14532 Stahnsdorf, Germany			
GIVEN NAME/FAMILY NAME Hans LEHRACH		INVENTOR'S SIGNATURE <i>Hans Lehrach</i>	DATE*
Residence (City, State & Country) Berlin, Germany		CITIZENSHIP Austrian	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country) Lutzelseiteinerweg 50, 14195 Berlin, Germany			
GIVEN NAME/FAMILY NAME		INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)		CITIZENSHIP	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			

I hereby appoint the following attorneys to prosecute this application and/or an international application based on this application and to transact all business in the Patent and Trademark Office connected therewith and in connection with the resulting patent based on instructions received from the entity who first sent the application papers to the attorneys identified below, unless the inventor(s) or assignee provides said attorneys with a written notice to the contrary:

Raymond C. Stewart	(Reg. No. 21,066)	Terrell C. Birch	(Reg. No. 19,382)
Joseph A. Kolasch	(Reg. No. 22,463)	James M. Slattery	(Reg. No. 28,380)
Bernard L. Sweeney	(Reg. No. 24,448)	Michael K. Mutter	(Reg. No. 29,680)
Charles Gorenstein	(Reg. No. 29,271)	Gerald M. Murphy, Jr.	(Reg. No. 28,977)
Leonard R. Svensson	(Reg. No. 30,330)	Terry L. Clark	(Reg. No. 32,644)
Andrew D. Meikle	(Reg. No. 32,868)	Marc S. Weiner	(Reg. No. 32,181)
Joe McKinney Muncy	(Reg. No. 32,334)	Donald J. Daley	(Reg. No. 34,313)
John W. Bailey	(Reg. No. 32,881)	John A. Castellano	(Reg. No. 35,094)
Gary D. Yacura	(Reg. No. 35,416)		

Send Correspondence to:

**BIRCH, STEWART, KOLASCH & BIRCH, LLP**

or **Customer No. 2292**

P.O. Box 747 • Falls Church, Virginia 22040-0747

Telephone: (703) 205-8000 • Facsimile: (703) 205-8050

PLEASE NOTE:

YOU MUST  
COMPLETE  
THE  
FOLLOWING:

Full Name of First  
Inventor  
Insert Name of  
Inventor  
Insert Date This  
Document is Signed

Insert Residence  
Insert Citizenship

Insert Post Office  
Address

Full Name of Second  
Inventor, if any  
see above

Full Name of Third  
Inventor, if any  
see above

Full Name of Fourth  
Inventor, if any  
see above

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

GIVEN NAME/FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Ivo Glynne GUT		02/10/80
Residence (City, State & Country)	CITIZENSHIP	
Paris, France	Swiss/British	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)		
18 rue du Moulin Vert, 75014 Paris, France		
GIVEN NAME/FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Kurt BERLIN		00/10/80
Residence (City, State & Country)	CITIZENSHIP	
Stahnsdorf, Germany	German	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)		
Marienkaferweg 4, 14532 Stahnsdorf, Germany		
GIVEN NAME/FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Hans LEHRACH		
Residence (City, State & Country)	CITIZENSHIP	
Berlin, Germany	Austrian	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)		
Lutzelsteinweg 50, 14195 Berlin, Germany		
GIVEN NAME/FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)	CITIZENSHIP	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)		

I hereby appoint the following attorneys to prosecute this application and/or an international application based on this application and to transact all business in the Patent and Trademark Office connected therewith and in connection with the resulting patent based on instructions received from the entity who first sent the application papers to the attorneys identified below, unless the inventor(s) or assignee provides said attorneys with a written notice to the contrary:

Raymond C. Stewart	(Reg. No. 21,066)	Terrell C. Birch	(Reg. No. 19,382)
Joseph A. Kolasch	(Reg. No. 22,463)	James M. Slattery	(Reg. No. 28,380)
Bernard L. Sweeney	(Reg. No. 24,448)	Michael K. Mutter	(Reg. No. 29,680)
Charles Gorenstein	(Reg. No. 29,271)	Gerald M. Murphy, Jr.	(Reg. No. 28,977)
Leonard R. Svensson	(Reg. No. 30,330)	Terry L. Clark	(Reg. No. 32,644)
Andrew D. Meikle	(Reg. No. 32,868)	Marc S. Weiner	(Reg. No. 32,181)
Joe McKinney Muncy	(Reg. No. 32,334)	Donald J. Daley	(Reg. No. 34,313)
John W. Bailey	(Reg. No. 32,881)	John A. Castellano	(Reg. No. 35,094)
Gary D. Yacura	(Reg. No. 35,416)		

Send Correspondence to:

**BIRCH, STEWART, KOLASCH & BIRCH, LLP**

or Customer No. 2292

P.O. Box 747 • Falls Church, Virginia 22040-0747

Telephone: (703) 205-8000 • Facsimile: (703) 205-8050

PLEASE NOTE:

YOU MUST

COMPLETE

THE

FOLLOWING:

↓

Full Name of First or Sole Inventor

Insert Name of Inventor

Insert Only This Document to Signed

→

Insert Residence

Insert Citizenship

→

Insert Post Office Address

→

Full Name of Second Inventor, if any

see above

Full Name of Third Inventor, if any

see above

Full Name of Fourth Inventor, if any

see above

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

GIVEN NAME/FAMILY NAME Ivo Glynne GUT		INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country) Paris, France		CITIZENSHIP Swiss/British	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country) 18 rue du Moulin Vert, 75014 Paris, France			
GIVEN NAME/FAMILY NAME Kurt BERLIN		INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country) Stahnsdorf, Germany		CITIZENSHIP German	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country) Marienkaferweg 4, 14532 Stahnsdorf, Germany			
GIVEN NAME/FAMILY NAME Hans LEHRACH		INVENTOR'S SIGNATURE	DATE* 1 Nov 2000
Residence (City, State & Country) Berlin, Germany		CITIZENSHIP Austrian	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country) <del>Kottbusser Platz 1, 10585 Berlin, Germany</del> Terrassenstrasse 31, 14129 Berlin, Germany			
GIVEN NAME/FAMILY NAME		INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)		CITIZENSHIP	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			

\*DATE OF SIGNATURE